

REPORT DOCUMENTATION PAGE					Form Approved OMB No. 0704-0188	
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</p>						
1. REPORT DATE (DD-MM-YYYY) 03/28/2017		2. REPORT TYPE Final			3. DATES COVERED (From - To) July 2015-December 2016	
4. TITLE AND SUBTITLE Neurophysiological outcomes of mTBI				5a. CONTRACT NUMBER		
				5b. GRANT NUMBER N00014-15-1-2799		
				5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S) Mark Tommerdahl, PhD (mark_tommerdahl@med.unc.edu)				5d. PROJECT NUMBER		
				5e. TASK NUMBER		
				5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of North Carolina at Chapel Hill Administrative Office Building, Suite 2200 104 Airport Drive, CB#1350 Chapel Hill, NC 27599-1350				8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 100 Alabama Street SW Suite 4R15 Atlanta GA 30303-3104				10. SPONSOR/MONITOR'S ACRONYM(S)		
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for Public Release; Distribution is Unlimited						
13. SUPPLEMENTARY NOTES						
14. ABSTRACT The proposed and executed effort of this project was a literature review of neurophysiological outcomes of mTBI. There are two sections in this report. The first section is an overview of neurophysiological outcomes of mTBI that have been addressed in the literature that are not related to cortical metrics. The second section addresses literature related to cortical metrics. Specifically, references of measures that can be routinely collected with the Brain Gauge (the device used to collect tactile based neurosensory assessments) were identified, collated, and excel spreadsheets were generated that describe the contents of these references.						
15. SUBJECT TERMS neurophysiological outcomes; mTBI; literature review; cortical metrics; Brain Gauge; tactile; neurosensory assessments; temporal order judgment; reaction time; timing perception						
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON	
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (Include area code)	
U	U	U	UU	21		

Neurophysiological outcomes of mTBI – Progress Report

The proposed and executed effort of this project was a literature review of neurophysiological outcomes of mTBI. There was no original research proposed or executed (i.e., no human subjects or animal research in this effort). There are two sections in this report. The first section (Part 1 - directly below) is an overview of neurophysiological outcomes of mTBI that have been addressed in the literature that are not related to cortical metrics. Subsequent sections address literature related to cortical metrics. Specifically, references of measures that can be routinely collected with the Brain Gauge (the device used to collect tactile based neurosensory assessments) were identified, collated and excel spreadsheets were generated that describe the contents of these references. These measures include temporal order judgement, reaction time, and timing perception and how those measures have been used for detecting deficits in information processing mechanisms in a number of cohorts, including mTBI.

Part 1.

A review of the literature on the topic of concussions shows what common sense should tell you—every concussion is different, so the physiology of concussions cannot be boiled down to a single, or even a single group, of statements. Physiological effects of concussion fall along a spectrum and are very much dependent on where on the head the person was hit, the force of the impact, the time since impact, total number of impacts the person has sustained, time between repeated impacts, age, and gender. Thus, data from the literature is varied and spans a wide range methods on study design and the subjects involved in testing.

Due to inconsistencies in subject injuries and time since impact, controlled animal studies can be of great help when evaluating physiological impacts of concussion. Animal studies, especially with rodents, allow for larger sample sizes, controlling the type and location of impact, investigating histological changes, and more standardized multiple impact studies. Homogenizing injuries and treatments allows for making more concrete statements about the impacts of concussion. For example, Huang et al. (2013) found greater tissue damage when impacts were administered 3d apart compared to 7d, and animals subjected to TBIs 3d apart still showed behavioral signs of concussion 1 month later while the other treatment group did not. On the other hand, concussions in humans are heterogeneous, with multiple sites of injury, torque, and confounding factors common due to the nature of injury. Thus looking at effects of a single impact at a single site does not necessarily reflect concussions that occur outside of a controlled environment. Furthermore, as reviewed by Shultz et al. (2016), there are many translational inconsistencies to consider when moving from animal models to humans, especially when using rodents. For example, because a rodent life span is vastly shorter than the average human, translating time between multiple impacts between species is not necessarily one-to-one.

Similarly, metabolism and pathophysiology differ greatly, making it important to interpret results with caution when relating to human TBI.

Balance is one measure commonly evaluated post-concussion, especially in athletics, although evidence and support as this being a reliable measure is limited. Studies have found that failure to pass the Balance Error Scoring System (BESS) test can be an effective measure for detecting concussions immediately following an impact (Guskiewicz et al. 2001, McCrea et al. 2005, Broglio & Puetz 2008), however false negatives are common—Guskiewicz et al. (2000) report only 30% of head injury patients show signs of balance problems immediately after injury. Furthermore, of those that do show alterations in balance, BESS scores usually return to equal or better than that of healthy controls within a week following the concussion, likely due to a practice effect and/or inter- and intra- rater variability (Riemann & Guskiewicz 2000, McCrea et al. 2003, Valovich et al. 2003, Vaolovich et al. 2004, Finnoff et al. 2009, Sheehan et al. 2011). Thus, reduced balance immediately after impact may be an effective way to quickly assess if a concussion has occurred, but not usually effective at determining when the injury has healed.

In contrast, imaging studies have shown altered brain function in the absence of diminished performance on cognitive tests in concussed individuals, but to date, no imaging study has been able to establish a significant link between continuing, chronic symptoms and physiological changes that result from concussion (reviewed in Shultz et al. 2016). While some concussions/TBI result in brain lesions, intracranial bleeding, or skull fractures detectable by CT or standard MRI, most are completely undetectable using these methods, especially when the injury is mild. Less than 10% of CT scans of concussion patients reveal any abnormality, and MRI abnormalities are present in only 10-57% of patients (reviewed in Bazarian et al. 2006). Keeping in mind that these studies included patients with severe TBI, the use of these standard imaging techniques appear to be very limited in their ability to diagnose concussion and mTBI.

Another MRI method capable of detecting physiological outcomes of concussion includes ASL, or arterial spin labeling, which can directly measure cerebral blood flow. This method has been used in relatively few studies to date to evaluate mTBI/concussion, with conflicting results found between studies. Wang et al. (2016) found decreased cerebral blood flow 24 hours after concussion in concussed athletes. After 8 day, when clinical symptoms had resolved, they documented that cerebral blood flow had decreased further from 24 h post-concussion levels, adding more evidence that the disappearance of clinical and neurocognitive symptoms commonly used to determine return to play is not necessarily an indicator that the athlete has fully recovered. However, Doshi et al. (2015) found that a week after concussion, athletes showed increased blood flow in the left striatum and frontal and occipital lobes compared to healthy controls. With small sample sizes (n=7 concussion group, n=12 control), this suggests that more studies are needed to determine the accuracy of this method.

Using fMRI pre- and post-concussion, Jantzen et al. (2004) found increased activation of the parietal and lateral frontal and cerebellar regions within 1 week of concussion, even in the absence of a change in cognitive performance postinjury. Similarly, McAllister et al. (1999) observed similar performance of concussed and control individuals, but there were differences in the activation patterns during working memory tasks between the two groups. Chen et al. (2004) also noted differences in working memory activation patterns despite normal performance in concussed individuals. Specifically, more activation was seen outside the area of interest that was not present in control groups, and BOLD (Blood Oxygen Level Dependent) responses were different between the two groups.

Changes in functional connectivity have been widely cited as a result of both acute and chronic concussion/mTBI. However, the details of such changes are not always clear, straightforward, or cohesive between studies depending on subjects used. Nevertheless, to make a generalization, it has been shown that even in asymptomatic patients, connectivity, as measured by resting state fMRI, is reduced post-concussion and is inversely related to the number of TBIs sustained by the individual (Johnson et al. 2012). In their review of the topic, Chong and Schwedt (2015) were only able to conclude that functional connectivity does appear to change following concussion, including alterations in the default-mode network (DMN), but they were not able to make any other broad generalizations on the topic.

Other techniques including EEG and DTI can also be effective at detecting altered brain function post-concussion. Again, results and conclusions are varied depending on severity of concussion and amount of time elapsed since impact. In general, DTI has documented changes in diffusivity both acutely and chronically (6 months post impact) (Henry et al. 2011), and Lipton et al. (2009) showed a correlation between executive function and activity in the dorsolateral prefrontal cortex with acute concussion cases (< 2 weeks post impact).

Another imaging method, PET, or more specifically [^{18}F]fluorodeoxyglucose (FDG)-PET, is able to detect changes in cortical metabolism post-concussion. As is the case with most concussion research, studies vary widely in severity and location of injury, and sample sizes are rarely big enough to make strong general conclusions. In their review of this topic, Byrnes et al. (2014) concluded that post-concussion, patients show a decrease in brain metabolism, either globally or regionally that can last days to months, with some indication of correlation with degree of injury.

Alterations in brain metabolite concentrations can be measured using magnetic resonance spectroscopy (MRS) as an indicator of disrupted metabolism within the brain (reviewed in Honce et al. 2016). This method has been used to show changes in creatine and glutamate/glutamine complex levels in asymptomatic high school football players following impact (Poole et al. 2014). Furthermore, the use of NAA/Cr and NAA/Cho ratios have been

successful in showing that athletes are not always fully recovered once clinical symptoms have resolved, and in athletes who sustain a second hit before metabolite ratios return to normal, recovery time from the second impact is significantly longer than in those who sustain a second hit after fully recovering from the first injury, as determined by MRS. Also, metabolite imbalances that were not present acutely can present up to 6 months post-injury (reviewed in Honce et al. 2016).

More recently, the use of blood biomarkers as an indicator of concussion has been more thoroughly investigated as a way to detect concussion more easily, cheaply, effectively, and without the radiation exposure of CT scans (reviewed in Zetterberg et al. 2013, Mondello et al. 2014, Plog & Nedergaard 2015, Huang et al. 2016, Kawata et al. 2016, Kulbe & Geddes 2016, Zetterberg & Blennow 2016). While they show more promise to accurately diagnosing mild TBI than imaging, they still have their downsides and there are currently no biomarkers tests available that can definitively detect mild concussions. In order to be effective, biomarkers must be present in detectable quantities in some body fluid, and they should be specific to CNS injury. As noted previously, each TBI patient is different and effects of impact are different in each patient depending on a number of factors. It is therefore difficult to find a single biomarker that can be used as a “gold standard” for TBI detection, particularly when considering that these molecules must 1) pass through the blood brain barrier (BBB) in detectable levels to be reliably and accurately measured in blood samples 2) measurement errors are possible at many levels including collection, storage, and measurement of samples 3) most of the biomarkers being investigated currently are not exclusive to head trauma and elevated levels can be an indicator of orthopedic or other peripheral injury. Analyzing CSF samples as opposed to blood can bypass some of these problems, but it is much more invasive and risky, and sampling cannot be repeated as often as a blood sample. That being said, there are some biomarkers that have been investigated and show some potential for helping to clarify the picture of what is going on inside a patient's head following impact. Specifically, S100- β , neuron-specific enolase (NSE), and glial fibrillary acidic protein (GFAP), are the most thoroughly studied biomarkers that show potential, and all have been extensively reviewed (Zetterberg et al. 2013, Mondello et al. 2014, Huang et al. 2016, Kawata et al. 2016, Kulbe & Geddes 2016, Pan et al. 2016, Zetterberg & Blennow 2016). While they show the more promise than other suggested biomarkers, they still have their downsides and have shown mixed results in their efficiency as an indicator of mTBI.

S100- β is a marker of activated astrocytes and is moved into the bloodstream by bulk flow via the glymphatic system (Plog et al 2015). However, it is also found in peripheral sources, including chondrocytes, adipocytes, and exocrine cells and elevated blood levels have been detected in patients following ischemic reperfusion injury, bone fractures, as well as in people with mood disorders (reviewed in Pan et al. 2016).

NSE is an enzyme involved in glycolysis, mostly within neurons, and it is upregulated in response to axon damage in order to maintain homeostasis. While multiple studies have shown significant increases blood levels of this protein following moderate and severe TBI, it has not been shown to be useful in detecting mild TBI. Furthermore, increased serum NSE has been detected following hypoperfusion, liver damage, kidney damage, migraines, and bone fractures, further limiting its ability to accurately detect TBI (Reviewed in Kawata et al. 2016).

GFAP is a structural protein found in astrocytes. With neurotrauma, astrocytes show morphological changes, including increases in size, which involves increased GFAP production, plus GFAP can be shed following astroglial damage either due to mechanical forces or necrosis (reviewed in Kawata et al. 2016). Unlike most other suggested biomarkers, this protein is not found outside the nervous system, making its presence in blood more specific to neurotrauma. Furthermore, multiple studies have shown that it can be found in detectable levels in mTBI patients, and levels appear to be correlated with severity of injury. This protein appears to hold more promise in detecting TBI, but it has not been studied as extensively as S100- β and many studies that have been done have had relatively small sample sizes. Additional studies are needed to verify its specificity and sensitivity, as well as establish how long it remains present in the blood (reviewed in Kawata et al. 2016, Kulbe & Geddes 2016, Pan et al. 2016).

With more testing and validation, using fluid biomarkers could be a valid part of determining when head trauma has occurred and/or when a patient has recovered. However, there are many confounding factors to consider and many shortcomings to this method, it has even been suggested that this method may never work to the degree that others have suggested due to the glymphatic system (Huang et al. 2016). In fact, Plog et al. (2015) found that serum levels of S100B, NSE, and GFAP were not elevated following experimentally induced TBI in mice when the glymphatic system was blocked. The potential for a false negative test result from any biomarkers measured from blood is a thus concern, particularly given that sleep deprivation (a possible side effect of TBI) as well as TBI in and of itself decreases glymphatic system activity (Iliff et al. 2014, Plog et al. 2015). Furthermore, even if proteins make it into the bloodstream, they will eventually degrade, be cleared by the hepatic and renal systems, and they can bind to carrier proteins, making standard measurement methods inaccurate (reviewed in Plog & Nedergaard 2015).

Huang et al. (2016) suggests a way around the obstacle of the glymphatic system by noting that TBI of any origin causes breakdown of the blood brain barrier (BBB). In theory, increased levels of circulating brain microvascular endothelial cells (cBMEC), the main constituent of the BBB, could be an indicator of TBI. But again, this suggestion has its drawbacks as increased cBMEC levels could be confounded by a number of other factors that also cause BBB breakdown, including (but not limited to) nicotine and HIV.

Recent advances in TBI detection have been in the field of molecular biology, using micro RNA (miRNA) as a biomarker for TBI. While studies in this area are limited, this method does appear to hold more promise than protein biomarkers discussed previously. Using miRNA as opposed to proteins allows for detection of the biomarker at lower concentrations, miRNA is less susceptible to degradation, both before and after sampling, evidence shows that there are more biomarkers that are specific to TBI and not influenced by peripheral damage, and there are miRNA markers that have been shown to be able to differentiate between mild, moderate, and severe TBI in mouse models and in humans (Balakathiresan et al. 2014, Sharma et al. 2014, Bhomia et al. 2016, Harrison et al. 2016, Sun et al. 2016, Taheri et al. 2016,).

Specifically, Sun et al. (2016), found that expression of the miRNA miR-23b was downregulated in plasma of TBI patients relative to healthy individuals. Bhomia et al. (2016), using blood samples from human subjects with either mTBI, severe TBI, orthopedic injury, or healthy controls, found 8 miRNAs specific to mTBI, 10 specific to sTBI, and 10 specific to orthopedic injury. In TBI patients, expression levels of most miRNAs showed a direct correlation to severity of injury as indicated by CT lesions. These results seem promising, but sample sizes were small (n=8/group) and all sampling was done 12-48 h after injury, meaning the ability to assess TBI status over time to track progress and recovery remains unknown. Further studies must also be done to determine if these miRNAs could also be present in other neurological conditions that may co-occur with TBI. For example, PTSD has been shown to have its own miRNA signature (Balakathiresan et al. 2016).

In summary, it has been widely documented that concussions/TBI cause a wide range of alterations in brain function. However, this injury is extremely heterogeneous in nature, making it difficult to make broad generalizations. Individual injuries can affect different parts of the brain in different ways depending on factors such as the location and intensity of impact, the number of previous concussions sustained, and time between impacts, so any method used to assess the injury must be able to do so irrespective of each of these factors. One generalization that can be made, however, is that changes in neurophysiology following concussion rarely are completely resolved at the same time that clinical symptoms disappear, and sustaining a second head injury before the first has resolved can drastically increase the severity of the injury. Thus, it is important to establish a method to accurately detect and track progress in order to prevent re-injury in an already injured brain.

References

Balakathiresan, NS, Chandran R, Bhomia M, Jia M, Li H, Maheshwari RK (2014). Serum and amygdala microRNA signatures of posttraumatic stress: Fear correlation and biomarker potential. *J Psychiatric Research* 57:65-73.

Bazarian JJ, Blyth B, Cimpello L (2006). Bench to bedside: Evidence for brain injury after concussion—looking beyond the computed tomography scan. *Academic Emergency Medicine* 13 (2):199-214.

Bhomia M, Balakathiresan NS, Wang KK, Papa L, Maheshwari RK (2016). A panel of serum miRNA biomarkers for the diagnosis of severe to mild traumatic brain injury in humans. *Scientific Reports* 6:28148.

Broglio SP & Puetz TW (2008). The effect of sport concussion on neurocognitive function, self-report symptoms and postural control; A meta-analysis. *Sports Med* 2008 38(1)53-67.

Byrnes KR, Wilson CM, Brabazon F, von Leden R, Jurgens JS, Oakes TR, Selwyn RG (2014). FDG-PET imaging in mild traumatic brain injury: a critical review. *Frontiers in Neuroimaging* 5(13).

Chen JK, Johnston KM, Frey S, Petrides M, Worsley K, Ptito A (2004). Functional abnormalities in symptomatic concussed athletes: an fMRI study. *NeuroImage* 22:68-82.

Chong CD and Schwedt TJ (2015). White matter damage and brain network alterations in concussed patients: a review of recent diffusion tensor imaging and resting-state functional connectivity data. *Curr Pain Headache Rep* 19(12):doi 10.1007/s11916-015-0485-0.

Doshi H, Wiseman N, Liu J, Wang W, Welch RD, O'Neil BJ, Zuk C, Wang X, Mika V, Szaflarski JP, Haacke EM, Kou Z (2015). Cerebral hemodynamic changes of mild traumatic brain injury at the acute stage. *PLoS One* 10:e0118061.

Finnoff JT, Peterson VJ, Hollman JH, Smith J (2009). Intrarater and interrater reliability of the Balance Error Scoring System (BESS). *American Academy of Physical Medicine and Rehabilitation* 1:50-54.

Guskiewicz KM, Ross SE, Marshall SW (2001). Postural stability and neuropsychological deficits after concussion in collegiate athletes. *Journal of Athletic Training* 36(3):263-273.

Guskiewicz KM, Weaver NL, Padua DA, Garrett WE (2000). Epidemiology of concussion in collegiate and high school football players. *Am J Sports Med* 28:643-650.

Harrison EB, Hochfelder CG, Lamberty BG, Meays BM, Morsey BM, Kelso ML, Fox HS, Yelamanchili SV (2016). Traumatic brain injury increases levels of miR-21 in extracellular vesicles: implications for neuroinflammation. *FEBS Open Bio*.

Henry LC, Tremblay J, Tremblay S, Lee A, Brun C, Lepore N, Theoret H, Ellemberg D, Lassonde M (2011). Acute and chronic changes in diffusivity measures after sports concussion. *Journal of Neurotrauma* 28:2049-2059.

Honce JM, Nyberg E, Jones I, Nagae L (2016). Neuroimaging of concussion. *Phys Med Rehabil Clin N Am* 27(2):411-428.

Huang L, Coats JS, Mohd-Yusof A, Yin Y, Assaad S, Muellner MJ, Kamper JE, Hartman RE, Dulcich M, Donovan VM, Oyoyo U, Obenaus A (2013). Tissue vulnerability is increased following repetitive mild traumatic brain injury in the rat. *Brain Research* 1499:109-120.

Huang S, Zhang B, Zhou Y (2016). Why molecular biomarkers of traumatic brain injury may never work: Effects of glymphatic pathway dysfunction. *Journal of Trauma and Treatment* 5(2): 309.

Iliff JJ, Chen MJ, Plog BA, Zeppenfeld DM, Soltero M, Yang L, Singh I, Deane R, Nedergaard M (2014). Impairment of glymphatic pathway function promotes tau pathology after traumatic brain injury. *The Journal of Neuroscience* 34(49):16180-16193.

Jantzen KJ, Anderson B, Steinberg FL, Scott Kelso JA (2004). A prospective functional MR imaging study of mild traumatic brain injury in college football players. *Am J Neuroradiol* 25:738-745.

Johnson B, Zhang K, Gay M, Horovitz S, Hallett M, Sebastianelli W, Slobounov S (2012). Alteration of brain default network in subacute phase of injury in concussed individuals: resting-state fMRI study. *Neuroimage* 59:511-518.

Kawata K, Liu CY, Merkel SF, Ramirez SH, Tierney RT, Langford D (2016). Blood biomarkers for brain injury: What are we measuring? *Neuroscience and Biobehavioral Reviews* 68:460-473.

Kulbe JR & Geddes JW (2016). Current status of fluid biomarkers in mild traumatic brain injury. *Experimental Neurology* 275:334-352.

Lipton ML, Gulko E, Zimmerman ME, Friedman BW, Kim M, Gellella E, Gold T, Shifteh K, Ardekani BA, Branch CA (2009). Diffusion-Tensor Imaging implicates prefrontal axonal injury in executive function impairment following very mild traumatic brain injury. *Neuroradiology* 252(3):816-824.

- McAllister TW, Saykin AJ, Flashman LA, Sparling MB, Johnson SC, Guerin SJ, Mamourian AC, Weaver JB, Yanofsky N (1999). Brain activation during working memory 1 month after traumatic brain injury: A functional MRI study. *Neurology* 53(6):1300-1308.
- McCrea M, Barr W, Guskiewicz K, Randolph C, Marshall SW, Cantu R, Onate JA, Kelly JP (2005). Standard regression-based methods for measuring recovery after sport-related concussion. *Journal of the International Neuropsychological Society* 11:58-69.
- McCrea M, Guskiewicz KM, Marshall SW, Barr W, Randolph C, Cantu RC, Onate JA, Yang J, Kelly JP (2003). Acute effects and recovery time following concussion in collegiate football players; The NCAA Concussion Study. *JAMA* 290(19): 2556-2563.
- Mondello S, Schmid K, Berger RP, Kobeissy F, Italiano D, Jeromin A, Hayes RL, Tortella FC, Buki A (2014). The challenge of mild traumatic brain injury: Role of biochemical markers in diagnosis of brain damage. *Medicinal Research Reviews* 34(3):503-31.
- Pan, J, Connolly ID, Dangelmajer S, Kintzing J, Ho AL, Grant G (2016). Sports-related brain injuries: connecting pathology to diagnosis. *Neurosurg Focus* 40(4):E14.
- Plog BA, Dashnaw ML, Hitomi E, Peng W, Liao Y, Lou N, Deane R, Nedergaard M (2015). Biomarkers of traumatic injury are transported from brain to blood via the glymphatic system. *The Journal of Neuroscience* 35(2):518-526.
- Plog BA & Nedergaard M (2015). Why have we not yet developed a simple blood test for TBI? *Expert Review of Neurotherapeutics* 15(5):465-468.
- Poole VN, Abbas K, Shenk TE, Breedlove EL, Breedlove KM, Robinson ME, Leverenz LJ, Nauman EA, Talavage TM, Dydak U (2014). MR Spectroscopic evidence of brain injury in the non-diagnosed collision sport athlete. *Developmental Neuropsychology* 39(6):459-473.
- Riemann BL & Guskiewicz KM (2000). Effects of mild head injury on postural stability as measured through clinical balance testing. *Journal of Athletic Training* 35(1):19-25.
- Shan R, Szmydynger-Chodobska J, Warren OU, Mohammad F, Zink BJ, Chodobski A (2016). A new panel of blood biomarkers for the diagnosis of mild traumatic brain injury/concussion in adults. *Journal of Neurotrauma* 33:49-57.
- Sharma A, Chandran R, Barry ES, Bhomia M, Hutchison MA, Balakathiresan NS, Grunberg NE, Maheshwari RK (2014). Identification of serum microRNA signatures for diagnosis of mild traumatic brain injury in a closed head injury model. *PLOS One* 9(11):e112019.

Sheehan DP & Lafave MR (2011). Intra-rater and inter-rater reliability of the Balance Error Scoring System in pre-adolescent school children. *Measurement in Physical Education and Exercise Science* 15:234-243.

Shultz SR, McDonald SJ, Vonder Haar C, Meconi A, Vink R, van Donkelaar P, Taneja C, Iverson GL, Christie BR (2016). The potential for animal models to provide insight into mild traumatic brain injury: Translational challenges and strategies. *Neuroscience and Biobehavioral Reviews* <http://dx.doi.org/10.1016/j.neubiorev.2016.09.014>

Sun L, Liu A, Zhang J, Ji W, Li Y, Yang X, Wu Z, Guo J (2016). miR-23b improves cognitive impairments in traumatic brain injury by targeting ATG12-mediated neuronal autophagy. *Behavioural Brain Research*.

Taheri S, Tanriverdi F, Zararsiz G, Elbuken G, Ulutabanca H, Karaca Z, Selcuklu A, Unluhizarci K, Tanriverdi K, Kelestimur F (2016). Circulating microRNAs as potential biomarkers for traumatic brain injury-induced hypopituitarism. *Journal of Neurotrauma*.

Valovich TC, Perrin DH, Gansneder BM (2003). Repeat administration elicits a practice effect with the Balance Error Scoring System but not with the standardized assessment of concussion in high school athletics. *Journal of Athletic Training* 38(1):51-56.

Valovich McLeod TC, Perrin DH, Guskiewicz KM, Shultz SJ, Diamond R, Gansneder BM (2004). Serial administration of clinical concussion assessments and learning effects in healthy young athletes. *Clin J Sport Med* 14(5):287-295.

Wang Y, Nelson LD, LaRoche AA, Pfaller AY, Nencka AS, Koch KM, McCrea MA (2016). Cerebral blood flow alterations in acute sports-related concussion. *Journal of Neurotrauma* 33:1227-1236.

Wright DK, Trezise J, Kamnaksh A, Bekdash R, Johnston LA, Ordidge R, Semple BD, Gardner AJ, Stanwell P, O'Brien TJ, Agoston DV, Shultz SR (2016). Behavioral, blood, and magnetic resonance imaging biomarkers of experimental mild traumatic brain injury. *Scientific Reports* 6:28713.

Zetterberg H, Smith DH, Blennow K (2013). Biomarkers of mild traumatic brain injury in cerebrospinal fluid and blood. *Nat Rev Neurol* 9:201-210.

Zetterberg H & Blennow K (2016). Fluid biomarkers for mild traumatic brain injury and related conditions. *Nature Reviews Neurology* 12:563-574.

Part 2.

Temporal order judgement (TOJ) defines an individual's ability to determine the order of two stimuli delivered to the peripheral sensory sheet. This ability is largely attributed to frontal-striatal pathways and TOJ is impacted in mTBI when that part of the cortex is impacted. Some neurological groups do particularly poorly in this task (e.g., schizophrenia and dyslexia). The contents of the TOJ references below are summarized in an excel spreadsheet.

Aoyama A, Krummenacher P, Palla A, Hilti LM, Brugger P. Impaired spatial-temporal integration of touch in xenomelia (body integrity identity disorder). *Spatial Cognition & Computation*. 2012 Apr 19;12(2-3):96-110.

Aglioti SM, Fiorio M, Forster B, Tinazzi M. Temporal discrimination of cross-modal and unimodal stimuli in generalized dystonia. *Neurology*. 2003 Mar 11;60(5):782-5.

Auld ML, Boyd R, Moseley GL, Ware R, Johnston LM. Tactile function in children with unilateral cerebral palsy compared to typically developing children. *Disability and rehabilitation*. 2012 Aug 1;34(17):1488-94.

Babkoff H, Zukerman GI, Fostick L, BEN-ARTZI EL. Effect of the diurnal rhythm and 24 h of sleep deprivation on dichotic temporal order judgment. *Journal of sleep research*. 2005 Mar 1;14(1):7-15.

Conte A, Khan N, Defazio G, Rothwell JC, Berardelli A. Pathophysiology of somatosensory abnormalities in Parkinson disease. *Nature Reviews Neurology*. 2013 Dec 1;9(12):687-97.

Craig JC, Rhodes RP, Busey TA, Kewley-Port D, Humes LE. Aging and tactile temporal order. *Attention, Perception, & Psychophysics*. 2010 Jan 1;72(1):226-35.

da Silva CF, Morgero KC, Mota AM, Piemonte ME, Baldo MV. Aging and Parkinson's disease as functional models of temporal order perception. *Neuropsychologia*. 2015 Nov 30;78:1-9.

Fiorio M, Tinazzi M, Bertolasi L, Aglioti SM. Temporal processing of visuotactile and tactile stimuli in writer's cramp. *Annals of neurology*. 2003 May 1;53(5):630-5.

Gibbon J, Rutschmann R. Temporal order judgment and reaction time. *Science*. 1969 Jul 25;165(3891):413-5.

Haenzi S, Stefanics G, Lanaras T, Calcagni M, Ghosh A. Botulinum toxin-A dose dependent perceptual loss on the hand after its cosmetic use on the face. *cortex*. 2014;63:118-20.

Heed T, Azañón E. Using time to investigate space: a review of tactile temporal order judgments as a window onto spatial processing in touch. *Frontiers in psychology*. 2014 Feb 17;5:76.

Kitazawa S, Moizumi S, Okuzumi A, Saito F, Shibuya S, Takahashi T, Wada M, Yamamoto S. Reversal of subjective temporal order due to sensory and motor integrations. *Sensorimotor foundations of higher cognition attention and performance*. 2008 Jan 30:73-97.

Laasonen M, Virsu VJ. Temporal order and processing acuity of visual, auditory, and tactile perception in developmentally dyslexic young adults. *Cognitive, Affective, & Behavioral Neuroscience*. 2001 Dec 1;1(4):394-410.

- Moseley GL, Gallace A, Spence C. Space-based, but not arm-based, shift in tactile processing in complex regional pain syndrome and its relationship to cooling of the affected limb. *Brain*. 2009 Nov 1;132(11):3142-51.
- Moseley GL, Gallagher L, Gallace A. Neglect-like tactile dysfunction in chronic back pain. *Neurology*. 2012 Jul 24;79(4):327-32.
- Nishikawa N, Shimo Y, Wada M, Hattori N, Kitazawa S. Effects of aging and idiopathic Parkinson's disease on tactile temporal order judgment. *PloS one*. 2015 Mar 11;10(3):e0118331.
- Pagel B, Heed T, Röder B. Change of reference frame for tactile localization during child development. *Developmental science*. 2009 Nov 1;12(6):929-37.
- Röder B, Rösler F, Spence C. Early vision impairs tactile perception in the blind. *Current Biology*. 2004 Jan 20;14(2):121-4.
- Sanger TD, Tarsy D, Pascual-Leone A. Abnormalities of spatial and temporal sensory discrimination in writer's cramp. *Movement disorders*. 2001 Jan 1;16(1):94-9.
- Sterr A, Green L, Elbert T. Blind Braille readers mislocate tactile stimuli. *Biological psychology*. 2003 May 31;63(2):117-27.
- Takahashi T, Kansaku K, Wada M, Shibuya S, Kitazawa S. Neural correlates of tactile temporal-order judgment in humans: an fMRI study. *Cerebral Cortex*. 2012 Jul 3:bhs179.
- Tinazzi M, Frasson E, Bertolasi L, Fiaschi A, Aglioti S. Temporal discrimination of somesthetic stimuli is impaired in dystonic patients. *Neuroreport*. 1999 May 14;10(7):1547-50.
- Tinazzi M, Fiaschi A, Frasson E, Fiorio M, Cortese F, Aglioti SM. Deficits of temporal discrimination in dystonia are independent from the spatial distance between the loci of tactile stimulation. *Movement disorders*. 2002 Mar 1;17(2):333-8.
- Tinazzi M, Fiorio M, Bertolasi L, Aglioti S. Timing of tactile and visuo-tactile events is impaired in patients with cervical dystonia. *Journal of neurology*. 2004 Jan 1;251(1):85-90.
- Van Damme S, Gallace A, Spence C, Crombez G, Moseley GL. Does the sight of physical threat induce a tactile processing bias?: Modality-specific attentional facilitation induced by viewing threatening pictures. *Brain research*. 2009 Feb 9;1253:100-6.
- Virsu V, Lahti-Nuuttila P, Laasonen M. Crossmodal temporal processing acuity impairment aggravates with age in developmental dyslexia. *Neuroscience letters*. 2003 Jan 23;336(3):151-4.
- Wada M, Yamamoto S, Kitazawa S. Effects of handedness on tactile temporal order judgment. *Neuropsychologia*. 2004 Dec 31;42(14):1887-95.
- Wada M, Suzuki M, Takaki A, Miyao M, Spence C, Kansaku K. Spatio-temporal processing of tactile stimuli in autistic children. *Scientific reports*. 2014 Aug 7;4:5985.

Reaction Time (RT). Reaction time describes an individual's information processing speed. The measure is simple in concept and easily measured, and for this reason, there are a large number of references on this topic that date back to 1885. However, the majority of commercial measures that collect reaction time introduce a very large variability for technical reasons and

tactile based reaction time is clearly the most reliable means of administering this very useful test. The contents of the RT references below are summarized in an excel spreadsheet.

Ancelin ML, Artero S, Portet F, Dupuy AM, Touchon J, Ritchie K. Non-degenerative mild cognitive impairment in elderly people and use of anticholinergic drugs: longitudinal cohort study. *Bmj*. 2006 Feb 23;332(7539):455-9.

Benton AL. Interactive effects of age and brain disease on reaction time. *Archives of Neurology*. 1977 Jun 1;34(6):369-70.

Caty G, Hu L, Legrain V, Plaghki L, Mouraux A. Psychophysical and electrophysiological evidence for nociceptive dysfunction in complex regional pain syndrome. *Pain®*. 2013 Nov 30;154(11):2521-8.

Davidson RJ, Leslie SC, Saron C. Reaction time measures of interhemispheric transfer time in reading disabled and normal children. *Neuropsychologia*. 1990 Dec 31;28(5):471-85.

Der G, Deary IJ. Age and sex differences in reaction time in adulthood: results from the United Kingdom Health and Lifestyle Survey. *Psychology and aging*. 2006 Mar;21(1):62.

Draper R, Manning A, O'Neill C. Two-signal discrimination: Relevance for psychopharmacology. *Drug Development Research*. 1982 Jan 1;2(S1):165-76.

Edwards AE, Cohen S. Visual illusion, tactile sensibility and reaction time under LSD-25. *Psychopharmacologia*. 1961 Sep 1;2(5):297-303.

Edwards L, Ring C, McIntyre D, Carroll D, Martin U. Psychomotor speed in hypertension: Effects of reaction time components, stimulus modality, and phase of the cardiac cycle. *Psychophysiology*. 2007 May 1;44(3):459-68.

Evarts EV, Teräsväinen H, Calne DB. Reaction time in Parkinson's disease. *Brain: a journal of neurology*. 1981 Mar;104(Pt 1):167-86.

Ferraro FR. No evidence of reaction time slowing in autism spectrum disorder. *Autism*. 2014 Dec 15;1362361314559986.

Fozard JL, Vercruyssen M, Reynolds SL, Hancock PA, Quilter RE. Age differences and changes in reaction time: the Baltimore Longitudinal Study of Aging. *Journal of gerontology*. 1994 Jul 1;49(4):P179-89.

Goodrich S, Henderson L, Kennard C. On the existence of an attention-demanding process peculiar to simple reaction time: converging evidence from Parkinson's disease. *Cognitive Neuropsychology*. 1989 May 1;6(3):309-31.

Hansson P, Ekblom A, Lindblom U, Marchettini P. Does acute intraoral pain alter cutaneous sensibility?. *Journal of Neurology, Neurosurgery & Psychiatry*. 1988 Aug 1;51(8):1032-6.

Hernández OH, Vogel-Sprott M, Huchín-Ramírez TC, Aké-Estrada F. Acute dose of alcohol affects cognitive components of reaction time to an omitted stimulus: differences among sensory systems. *Psychopharmacology*. 2006 Jan 1;184(1):75-81.

Hick WE. On the rate of gain of information. *Quarterly Journal of Experimental Psychology*. 1952 Mar 1;4(1):11-26.

Klemmer ET. Simple reaction time as a function of time uncertainty. *Journal of experimental psychology*. 1957 Sep;54(3):195.

Lajoie Y, Gallagher S. Predicting falls within the elderly community: comparison of postural sway, reaction time, the Berg balance scale and the Activities-specific Balance Confidence (ABC) scale for comparing fallers and non-fallers. *Archives of gerontology and geriatrics*. 2004 Feb 29;38(1):11-26.

Leonard JA. Tactual choice reactions: I. *Quarterly Journal of Experimental Psychology*. 1959 May 1;11(2):76-83.

Lord SR, Clark RD. Simple physiological and clinical tests for the accurate prediction of falling in older people. *Gerontology*. 1996 Jul 1;42(4):199-203.

Lorenzo I, Ramos J, Arce C, Guevara MA, Corsi-Cabrera M. Effect of total sleep deprivation on reaction time and waking EEG activity in man. *SLEEP-NEW YORK*. 1995 Jun 1;18:346-.

MacDonald SW, Nyberg L, Bäckman L. Intra-individual variability in behavior: links to brain structure, neurotransmission and neuronal activity. *Trends in neurosciences*. 2006 Aug 31;29(8):474-80.

MacFlynn G, Montgomery EA, Fenton GW, Rutherford W. Measurement of reaction time following minor head injury. *Journal of Neurology, Neurosurgery & Psychiatry*. 1984 Dec 1;47(12):1326-31.

Meere J, Vreeling HJ, Sergeant J. A Motor Presetting Dstudy in Hyperactive, Learning Disabled and Control Children. *Journal of Child Psychology and Psychiatry*. 1992 Nov 1;33(8):1347-51.

Merkel J. Die zeitlichen verhältnisse der willensthatigkeit. *Philos Stud*. 2:73-127.

Moreau V, Berquin AD, Plaghki L. Laser-evoked potentials correlate with clinical evolution in a case of spontaneous and recurrent complex regional pain syndrome type I. *The Clinical journal of pain*. 2007 May 1;23(4):375-9.

Patil S, Phatale S. Auditory and Visual Reaction Time-A Tool for Early Detection of Neuropathy in Diabetics. *International Journal of Health Sciences and Research (IJHSR)*. 2015;5(4):141-6.

Ponsford J, Kinsella G. Attentional deficits following closed-head injury. *Journal of Clinical and Experimental Neuropsychology*. 1992 Sep 1;14(5):822-38.

Puts NA, Wodka EL, Tommerdahl M, Mostofsky SH, Edden RA. Impaired tactile processing in children with autism spectrum disorder. *Journal of Neurophysiology*. 2014 May 1;111(9):1803-11.

Reinersmann A, Haarmeyer GS, Blankenburg M, Frettlöh J, Krumova EK, Ocklenburg S, Maier C. Left is where the L is right. Significantly delayed reaction time in limb laterality recognition in both CRPS and phantom limb pain patients. *Neuroscience letters*. 2010 Dec 17;486(3):240-5.

Richerson SJ, Robinson CJ, Shum J. A comparative study of reaction times between type II diabetics and non-diabetics. *Biomedical engineering online*. 2005 Feb 21;4(1):12.

Ruesch J. Dark adaptation, negative after images, tachistoscopic examinations and reaction time in head injuries. *Journal of Neurosurgery*. 1944 Jul;1(4):243-51.

Sarno S, Erasmus LP, Lipp B, Schlaegel W. Multisensory integration after traumatic brain injury: a reaction time study between pairings of vision, touch and audition. *Brain Injury*. 2003 Jan 1;17(5):413-26.

Schwartz F, Carr AC, Munich RL, Glauber S, Lesser B, Murray J. Reaction time impairment in schizophrenia and affective illness: the role of attention. *Biological Psychiatry*. 1989 Mar 1;25(5):540-8.

Sherwood DE, Selder DJ. Cardiorespiratory health, reaction time and aging. *Medicine and science in sports*. 1978 Dec;11(2):186-9.

Shucard DW, Benedict RH, Tekok-Kilic A, Lichter DG. Slowed reaction time during a continuous performance test in children with Tourette's syndrome. *Neuropsychology*. 1997 Jan;11(1):147.

Spence C, Lloyd D, McGlone F, Nicholls ME, Driver J. Inhibition of return is supramodal: a demonstration between all possible pairings of vision, touch, and audition. *Experimental Brain Research*. 2000 Sep 1;134(1):42-8.

Stuss DT, Stethem LL, Hugenholtz H, Picton T, Pivik J, Richard MT. Reaction time after head injury: fatigue, divided and focused attention, and consistency of performance. *Journal of Neurology, Neurosurgery & Psychiatry*. 1989 Jun 1;52(6):742-8.

Tamm L, Narad ME, Antonini TN, O'Brien KM, Hawk LW, Epstein JN. Reaction time variability in ADHD: a review. *Neurotherapeutics*. 2012 Jul 1;9(3):500-8.

ten Hoopen G, Akerboom S, Raaymakers E. Vibrotactile choice reaction time, tactile receptor systems and ideomotor compatibility. *Acta Psychologica*. 1982 Mar 31;50(2):143-57.

Zahn TP, Mirsky AF. Reaction time indicators of attention deficits in closed head injury. *Journal of Clinical and Experimental Neuropsychology*. 1999 Jun 1;21(3):352-67.

Timing Perception. Timing perception is an individual's ability to accurately perceive the duration of a stimulus. Although this is reliant on a number of nodes in the neural network, the cerebellum plays a large role in an individual's ability to differentiate between two stimuli that differ only in duration. . The contents of the timing perception references below are summarized in an excel spreadsheet.

Anderson JW, Schmitter-Edgecombe M. Recovery of time estimation following moderate to severe traumatic brain injury. *Neuropsychology*. 2011 Jan;25(1):36.

Brannon EM, Roussel LW, Meck WH, Woldorff M. Timing in the baby brain. *Cognitive Brain Research*. 2004 Oct 31;21(2):227-33.

Carrasco MC, Redolat R, Simon VM. Effects of cigarette smoking on time estimation. *Human Psychopharmacology: Clinical and Experimental*. 1998 Dec 1;13(8):565-73.

Carmen Carrasco, M. Jesus Guillem, Rosa Redolat M. Estimation of short temporal intervals in Alzheimer's disease. *Experimental Aging Research*. 2000 Apr 1;26(2):139-51.

Carrasco MC, Bernal MC, Redolat R. Time estimation and aging: a comparison between young and elderly adults. *The International Journal of Aging and Human Development*. 2001 Mar;52(2):91-101.

Carroll CA, O'Donnell BF, Shekhar A, Hetrick WP. Timing dysfunctions in schizophrenia span from millisecond to several-second durations. *Brain and cognition*. 2009 Jul 31;70(2):181-90.

Casini L, Ramdani-Beauvir C, Burle B, Vidal F. How does one night of sleep deprivation affect the internal clock?. *Neuropsychologia*. 2013 Jan 31;51(2):275-83.

Davalos DB, Kisley MA, Freedman R. Behavioral and electrophysiological indices of temporal processing dysfunction in schizophrenia. *The Journal of neuropsychiatry and clinical neurosciences*. 2005 Nov;17(4):517-25.

Davalos DB, Kisley MA, Ross RG. Deficits in auditory and visual temporal perception in schizophrenia. *Cognitive Neuropsychiatry*. 2002 Nov 1;7(4):273-82.

Davalos DB, Kisley MA, Ross RG. Effects of interval duration on temporal processing in schizophrenia. *Brain and cognition*. 2003 Aug 31;52(3):295-301.

Eagleman DM. Using time perception to measure fitness for duty. *Military Psychology*. 2009;21(S1):S123.

Elvevåg B, McCormack T, Gilbert A, Brown GD, Weinberger DR, Goldberg TE. Duration judgements in patients with schizophrenia. *Psychological medicine*. 2003 Nov 1;33(07):1249-61.

Elvevåg B, Brown GD, McCormack T, Vousden JI, Goldberg TE. Identification of tone duration, line length, and letter position: an experimental approach to timing and working memory deficits in schizophrenia. *Journal of abnormal psychology*. 2004 Nov;113(4):509.

Gilaie-Dotan S, Ashkenazi H, Dar R. A Possible Link between Supra-Second Open-Ended Timing Sensitivity and Obsessive-Compulsive Tendencies. *Frontiers in Behavioral Neuroscience*. 2016;10.

Gilden DL, Marusich LR. Contraction of time in attention-deficit hyperactivity disorder. *Neuropsychology*. 2009 Mar;23(2):265.

Gooch D, Snowling M, Hulme C. Time perception, phonological skills and executive function in children with dyslexia and/or ADHD symptoms. *Journal of Child Psychology and Psychiatry*. 2011 Feb 1;52(2):195-203.

Graham-Schmidt KT, Martin-Iverson MT, Holmes NP, Waters FA. When one's sense of agency goes wrong: Absent modulation of time perception by voluntary actions and reduction of perceived length of intervals in passivity symptoms in schizophrenia. *Consciousness and Cognition*. 2016 Oct 31;45:9-23.

Grondin S, Rousseau R. Judging the relative duration of multimodal short empty time intervals. *Attention, Perception, & Psychophysics*. 1991 May 1;49(3):245-56.

Grondin S. Timing and time perception: a review of recent behavioral and neuroscience findings and theoretical directions. *Attention, Perception, & Psychophysics*. 2010 Apr 1;72(3):561-82.

Grondin S, Kuroda T, Mitsudo T. Spatial effects on tactile duration categorization. *Canadian Journal of Experimental Psychology/Revue canadienne de psychologie expérimentale*. 2011 Sep;65(3):163.

Gruber RP, Block RA. Effect of caffeine on prospective and retrospective duration judgements. *Human Psychopharmacology: Clinical and Experimental*. 2003 Jul 1;18(5):351-9.

Güçlü B, Sevinc E, Canbeyli R. Duration discrimination by musicians and nonmusicians. *Psychological reports*. 2011 Jun;108(3):675-87.

Halberstadt AL, Sindhunata IS, Scheffers K, Flynn AD, Sharp RF, Geyer MA, Young JW. Effect of 5-HT 2A and 5-HT 2C receptors on temporal discrimination by mice. *Neuropharmacology*. 2016 Aug 31;107:364-75.

Hinton SC, Meck WH. Increasing the speed of an internal clock: the effects of nicotine on interval timing. *Drug development research*. 1996 Jul 1;38(3-4):204-11.

- Jones LA, Poliakoff E, Wells J. Good vibrations: Human interval timing in the vibrotactile modality. *The Quarterly Journal of Experimental Psychology*. 2009 Nov 1;62(11):2171-86.
- Koch G, Oliveri M, Caltagirone C. Neural networks engaged in milliseconds and seconds time processing: evidence from transcranial magnetic stimulation and patients with cortical or subcortical dysfunction. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*. 2009 Jul 12;364(1525):1907-18.
- Lee KH, Dixon JK, Spence SA, Woodruff PW. Time perception dysfunction in psychometric schizotypy. *Personality and individual differences*. 2006 May 31;40(7):1363-73.
- Lee KH, Bhaker RS, Mysore A, Parks RW, Birkett PB, Woodruff PW. Time perception and its neuropsychological correlates in patients with schizophrenia and in healthy volunteers. *Psychiatry research*. 2009 Apr 30;166(2):174-83.
- Maricq AV, Church RM. The differential effects of haloperidol and methamphetamine on time estimation in the rat. *Psychopharmacology*. 1983 Jan 1;79(1):10-5.
- McDonald J, Schleifer L, Richards JB, de Wit H. Effects of THC on behavioral measures of impulsivity in humans. *Neuropsychopharmacology*. 2003 Jul 1;28(7):1356.
- Meck WH. Neuropharmacology of timing and time perception. *Cognitive brain research*. 1996 Jun 30;3(3):227-42.
- Meyers CA, Levin HS. Temporal perception following closed head injury: Relationship of orientation and attention span. *Cognitive and Behavioral Neurology*. 1992 Jan 1;5(1):28-32.
- Mioni G, Stablum F, McClintock SM, Cantagallo A. Time-based prospective memory in severe traumatic brain injury patients: The involvement of executive functions and time perception. *Journal of the International Neuropsychological Society*. 2012 Jul 1;18(04):697-705.
- Mioni G, Stablum F, Cantagallo A. Time discrimination in traumatic brain injury patients. *Journal of clinical and experimental neuropsychology*. 2013 Jan 1;35(1):90-102.
- Mioni G, Stablum F, Prunetti E, Grondin S. Time perception in anxious and depressed patients: A comparison between time reproduction and time production tasks. *Journal of affective disorders*. 2016 May 15;196:154-63.
- Perbal S, Couillet J, Azouvi P, Pouthas V. Relationships between time estimation, memory, attention, and processing speed in patients with severe traumatic brain injury. *Neuropsychologia*. 2003 Dec 31;41(12):1599-610.
- Radonovich KJ, Mostofsky SH. Duration judgments in children with ADHD suggest deficient utilization of temporal information rather than general impairment in timing. *Child Neuropsychology*. 2004 Sep 1;10(3):162-72.
- Rammsayer TH. Neuropharmacological evidence for different timing mechanisms in humans. *The Quarterly Journal of Experimental Psychology: Section B*. 1999 Aug 1;52(3):273-86.
- Rammsayer TH. The effects of sensory modality and type of task on discrimination of durations ranging from 400 to 1,400 milliseconds. *Proceedings of Fechner Day*. 2010;26(1):439-44.
- Righi S, Gronchi G, Paganini M, Piacentini S, Viggiano MP. Huntington's Disease and Scalar Expectancy Theory: A Memory-Based Time Perception Deficit. InEAPCogSci 2015.

Rubia K, Taylor A, Taylor E, Sergeant JA. Synchronization, anticipation, and consistency in motor timing of children with dimensionally defined attention deficit hyperactivity behaviour. *Perceptual and motor skills*. 1999 Dec;89(3_suppl):1237-58.

Rubia K, Noorloos J, Smith A, Gunning B, Sergeant J. Motor timing deficits in community and clinical boys with hyperactive behavior: the effect of methylphenidate on motor timing. *Journal of abnormal child psychology*. 2003 Jun 1;31(3):301-13.

Schmitter-Edgecombe M, Rueda AD. Time estimation and episodic memory following traumatic brain injury. *Journal of Clinical and Experimental Neuropsychology*. 2008 Jan 31;30(2):212-23.

Sewell RA, Schnakenberg A, Elander J, Radhakrishnan R, Williams A, Skosnik PD, Pittman B, Ranganathan M, D'Souza DC. Acute effects of THC on time perception in frequent and infrequent cannabis users. *Psychopharmacology*. 2013 Mar 1;226(2):401-13.

Shebloski KL, Broadway JM. Commentary: Effects of psilocybin on time perception and temporal control of behavior in humans. *Frontiers in Psychology*. 2016;7:736. doi:10.3389/fpsyg.2016.00736.

Sil'kis IG. Possible mechanisms for the effects of neuromodulators on the perception of time intervals. *Neurochemical Journal*. 2012 Apr 1;6(2):144-52.

Smith A, Taylor E, Warner Rogers J, Newman S, Rubia K. Evidence for a pure time perception deficit in children with ADHD. *Journal of Child Psychology and Psychiatry*. 2002 May 1;43(4):529-42.

Smith AB, Taylor E, Brammer M, Halari R, Rubia K. Reduced activation in right lateral prefrontal cortex and anterior cingulate gyrus in medication-naïve adolescents with attention deficit hyperactivity disorder during time discrimination. *Journal of Child Psychology and Psychiatry*. 2008 Sep 1;49(9):977-85.

Smith JG, Harper DN, Gittings D, Abernethy D. The effect of Parkinson's disease on time estimation as a function of stimulus duration range and modality. *Brain and cognition*. 2007 Jul 31;64(2):130-43.

Teixeira S, Machado S, Paes F, Velasques B, Guilherme Silva J, L Sanfim A, Minc D, Anghinah R, L Menegaldo L, Salama M, Cagy M. Time perception distortion in neuropsychiatric and neurological disorders. *CNS & Neurological Disorders-Drug Targets (Formerly Current Drug Targets-CNS & Neurological Disorders)*. 2013 Aug 1;12(5):567-82.

Toplak ME, Tannock R. Time perception: modality and duration effects in attention-deficit/hyperactivity disorder (ADHD). *Journal of abnormal child psychology*. 2005 Oct 1;33(5):639-54.

Toplak ME, Rucklidge JJ, Hetherington R, John SC, Tannock R. Time perception deficits in attention-deficit/hyperactivity disorder and comorbid reading difficulties in child and adolescent samples. *Journal of Child Psychology and Psychiatry*. 2003 Sep 1;44(6):888-903.

Toplak ME, Dockstader C, Tannock R. Temporal information processing in ADHD: findings to date and new methods. *Journal of neuroscience methods*. 2006 Feb 15;151(1):15-29.

Turgeon M, Lustig C, Meck WH. Cognitive aging and time perception: roles of Bayesian optimization and degeneracy. *Frontiers in aging neuroscience*. 2016;8.

Van der Lubbe RH, Van Mierlo CM, Postma A. The involvement of occipital cortex in the early blind in auditory and tactile duration discrimination tasks. *Journal of Cognitive Neuroscience*. 2010 Jul;22(7):1541-56.

VanMarle K, Wynn K. Six-month-old infants use analog magnitudes to represent duration. *Developmental science*. 2006 Sep 1;9(5).

Wearden JH, Lejeune H. Scalar properties in human timing: Conformity and violations. *The Quarterly Journal of Experimental Psychology*. 2008 Apr 1;61(4):569-87.

Wearden JH, Smith-Spark JH, Cousins R, Edelstyn NM, Cody FW, O'Boyle DJ. Stimulus timing by people with Parkinson's disease. *Brain and cognition*. 2008 Aug 31;67(3):264-79.

Westheimer G. Discrimination of short time intervals by the human observer. *Experimental Brain Research*. 1999 Oct 1;129(1):121-6.

Wittmann M, Carter O, Hasler F, Cahn BR, Grimberg U, Spring P, Hell D, Flohr H, Vollenweider FX. Effects of psilocybin on time perception and temporal control of behaviour in humans. *Journal of Psychopharmacology*. 2007 Jan;21(1):50-64.

Frequency discrimination. Frequency discrimination is an individuals' ability to differentiate between two sinusoidal stimuli on the basis of frequency. This has been of ongoing interest in the somatosensory research community for decades, but very few have a clue as to what frequency discrimination relies on. Although it is fairly evident that frequency discrimination relies on synchronization of cortical ensembles, a number of researchers have designed experiments to push an alternative hypothesis (by designing out that possible conclusion). Some of the references pertaining to frequency discrimination are listed below and the contents of these are summarized in an excel spreadsheet.

Flor H, Denke C, Schaefer M, Grüsser S. Effect of sensory discrimination training on cortical reorganisation and phantom limb pain. *The Lancet*. 2001 Jun 2;357(9270):1763-4.

Hernández A, Zainos A, Romo R. Neuronal correlates of sensory discrimination in the somatosensory cortex. *Proceedings of the National Academy of Sciences*. 2000 May 23;97(11):6191-6.

Hollins M, Sigurdsson A. Vibrotactile amplitude and frequency discrimination in temporomandibular disorders. *Pain*. 1998 Mar 31;75(1):59-67.

Levänen S, Hamdorf D. Feeling vibrations: enhanced tactile sensitivity in congenitally deaf humans. *Neuroscience letters*. 2001 Mar 23;301(1):75-7.

Luna R, Hernández A, Brody CD, Romo R. Neural codes for perceptual discrimination in primary somatosensory cortex. *Nature neuroscience*. 2005 Sep 1;8(9):1210-9.

Mahns DA, Perkins NM, Sahai V, Robinson L, Rowe MJ. Vibrotactile frequency discrimination in human hairy skin. *Journal of neurophysiology*. 2006 Mar 1;95(3):1442-50.

Recanzone GH, Merzenich MM, Jenkins WM, Grajski KA, Dinse HR. Topographic reorganization of the hand representation in cortical area 3b owl monkeys trained in a frequency-discrimination task. *Journal of neurophysiology*. 1992 May 1;67(5):1031-56.

Recanzone GH, Jenkins WM, Hradek GT, Merzenich MM. Progressive improvement in discriminative abilities in adult owl monkeys performing a tactile frequency discrimination task. *Journal of Neurophysiology*. 1992 May 1;67(5):1015-30.

Recanzone GH, Merzenich MM, Jenkins WM. Frequency discrimination training engaging a restricted skin surface results in an emergence of a cutaneous response zone in cortical area 3a. *Journal of Neurophysiology*. 1992 May 1;67(5):1057-70.

Sörös P, Marmurek J, Tam F, Baker N, Staines WR, Graham SJ. Functional MRI of working memory and selective attention in vibrotactile frequency discrimination. *BMC neuroscience*. 2007 Jul 4;8(1):48.

Stuart M, Turman AB, Shaw J, Walsh N, Nguyen V. Effects of aging on vibration detection thresholds at various body regions. *BMC geriatrics*. 2003 Feb 25;3(1):1.